




# *Hypena madefactalis* Gray-edged Bomolocha



FAMILY: Erebidae SUBFAMILY: Hypeninae TRIBE:

TAXONOMIC COMMENTS: One of 29 species in this genus that occur in North America north of Mexico (Lafontaine and Schmidt, 2010). Fifteen species have been recorded in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984; as *Bomolocha madefactalis*); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: North Carolina records come from a variety of rich hardwood forest, ranging from mesic, high elevation stands, cove forests, and Amphibolite ridges in the Mountains and alluvial and mafic forests in the Piedmont. Coastal Plain sites may all be associated with planted walnuts.

FOOD: Larvae feed on Black Walnut; one was also reared on Shagbark Hickory (Wagner et al., 2011)

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S4S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: